

BLOCH'S THEOREM FOR ALGEBROID MULTIFUNCTIONS II

DOMINIQUE GUILLOT AND THOMAS RANSFORD

ABSTRACT. Bloch's theorem is extended to multifunctions of the form

$$F(z) = \{w \in \mathbb{C} : w^n + a_1(z)w^{n-1} + \cdots + a_{n-1}(z)w + a_n(z) = 0\} \quad (|z| < 1),$$

where $a_1(z), \dots, a_n(z)$ are holomorphic functions on the unit disk and $a_1'(0) = 1$.

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ LAVAL, QUÉBEC (QC), CANADA G1K 7P4
E-mail address: `dguillot@mat.ulaval.ca`

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ LAVAL, QUÉBEC (QC), CANADA G1K 7P4
E-mail address: `ransford@mat.ulaval.ca`

2000 *Mathematics Subject Classification.* Primary 30C25, Secondary 30G30, 32A12.

Key words and phrases. Bloch's theorem, Landau's constant, multifunction, normal family.

First author partially supported by an NSERC undergraduate student research award.

Second author partially supported by grants from NSERC and the Canada research chairs program.