

RANDOM PROJECTIONS AND GOODNESS-OF-FIT TESTS IN INFINITE-DIMENSIONAL SPACES

JUAN ANTONIO CUESTA-ALBERTOS, RICARDO FRAIMAN, AND THOMAS RANSFORD

ABSTRACT. In this paper we provide conditions under which a distribution is determined by just one randomly chosen distribution. Then we apply our results to construct goodness-of-fit tests for the one- and two-sample problems. We include some simulations as well as the application of our results to a real data set.

Our results are valid for every separable Hilbert space.

DEPARTAMENTO DE MATEMÁTICAS, ESTADÍSTICA Y COMPUTACIÓN, UNIVERSIDAD DE CANTABRIA, SPAIN
E-mail address: `cuestaj@unican.es`

DEPARTAMENTO DE MATEMÁTICA Y CIENCIAS, UNIVERSIDAD DE SAN ANDRÉS, ARGENTINA
E-mail address: `rfraiman@udesa.edu.ar`

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ LAVAL, QUÉBEC (QC), CANADA G1K 7P4
E-mail address: `ransford@mat.ulaval.ca`

2000 *Mathematics Subject Classification.* Primary 62H15; Secondary 28C20, 60B11 and 60E05.

Key words and phrases. Cramér-Wold theorem, random projections, Hilbert spaces, goodness-of-fit tests, Kolmogorov-Smirnov projected test, single null hypothesis, two samples.

First author partially supported by the Spanish Ministerio de Ciencia y Tecnología, grant MTM2005-08519-C02-02.

Third author partially supported by grants from NSERC and the Canada research chairs program.